

The prevalence of lower respiratory tract infection in adults visiting Tribhuvan University Teaching Hospital

P. Gauchan, B. Lekhak, J. B. Sherchand

Tribhuvan University Institute of Medicine, Department of Clinical Microbiology and Health Research Laboratory, Central Department of Microbiology, Tribhuvan University, Kirtipur.

Correspondence to: Punita Gauchan, Department of Clinical Microbiology and Health Research Laboratory, Tribhuvan University, Institute of Medicine, Kathmandu, Nepal
e-mail: punita.gauchan@gmail.com

Background: Respiratory conditions impose enormous burden on society. Reports indicated that the top five respiratory diseases accounted for 17.4% of all deaths and 13.3% of all Disability-Adjusted Life Years (DALYs). Also, out of total acute respiratory disease, 20-24% of deaths are accounted for by Lower Respiratory Tract Infection (LRTI). In developing countries like Nepal the need for timely diagnosis of the cases and the administration of appropriate therapy based on the antibiotic susceptibility test of the causative agents is critical. However, emergence of resistant strains may occur during antibiotic therapy, which is one of the contributing factors for the increase in the frequency of LRTI in recent years in the adult population of Nepal as well.

Methods: A hospital based cross-sectional study was carried out from March 2002 to February 2003. Total 181 adults presenting with LRTI defined by a new or increasing cough, productive sputum, chest pain, fever, anorexia, haemoptysis, headaches and throat ache were enrolled with their consent. This is a prospective study which included bacteriological culture, microscopic examination and sensitivity testing of bacterial isolates *in vitro* in Health Research laboratory following Standard Operating Procedures (SOPs).

Results: Lower Respiratory Infection was found prevalent in 75 cases (41.4%). Males (61.3%) were found more at risk to LRTI than females (38.7%). LRTI was found most prevalent in 50-59 year age groups (21.3%). Altogether 15 different types of bacteria were identified majority of which were Gram-negative bacteria (72.4%). *Haemophilus influenzae* was the commonest isolate at 23.0% followed by *Klebsiella pneumoniae* (18.3%). Among Gram-positive isolates *Streptococcus pneumoniae* was predominant (12.7%) followed by *Staphylococcus aureus* (9.3%). The *in vitro* antibiotic susceptibility test of the isolates showed that Chloramphenicol (100%) was the most effective antibiotic against Gram-negative bacteria followed by Amikacin (79.1%) and Ciprofloxacin (66.7%), and the least effective was Co-trimoxazole (20.6%). Similarly, for the Gram-positive bacteria Ciprofloxacin (79.2%) was the most effective antibiotic and the least effective was Co-trimoxazole.

Conclusion: The study shows increasing number of respiratory pathogens resistant to antimicrobials in-use to treat the infection.

Key words: LRTI, Bacteria, Antibiotic susceptibility pattern, Respiratory pathogen.

Lower respiratory tract infection in adults

Introduction

Respiratory diseases were responsible for 1/5 of youth and adult deaths in 1998. These deaths were mainly caused by Chronic Obstructive Pulmonary Disease (COPD), Tuberculosis (TB), and Acute Respiratory Infections (ARI; mostly pneumonia). Among infectious diseases, ARI and Tuberculosis were two of the six leading causes of death across all ages.¹

The causes of the community-acquired Lower Respiratory Tract Infection (LRTI) are not well recognized, although it is traditionally taught that most are caused by viruses and atypical pathogens. There is a growing need for the better understanding of the pathogens that cause these infections which should allow a logical approach to treatment. The aetiology and the symptomatology of respiratory diseases vary with age, season, the type of population at risk, and other factors.² The need for the timely diagnosis of the cases and the administration of appropriate therapy based on the antibiotic susceptibility test of the causative agent is critical for blocking further spread of the pathogen which might otherwise lead to complications. Still, resistant strains may emerge during antibiotic therapy³ which is one of the contributing factors for the increase in the frequency of LRTI in recent years in the adult population of Nepal as well.

In view to obtain information regarding prevalence of LRTI, aetiology as well as to have knowledge in the current antimicrobials susceptibility pattern of the pathogens, this study was done.

Material and Methods

This was a hospital based prospective study done on 181 patients visiting out patient department of Tribhuvan University Teaching Hospital. Sputum for bacteriological culture and parasitological and fungal examination were collected after informed consent from patients presenting with LRTI as defined by a new or increasing cough, productive sputum, chest pain, fever, anorexia, haemoptysis, headache and weight loss. Before culture Gram-stained smear of every specimen was first examined microscopically. In microscopic examination sputum showing less than 10 squamous epithelial cells and more than 25 leucocytes or pus cells per low power field confirmed the reliability of the specimen indicating that it was not contaminated with saliva. Pathogens were isolated using Mac Conkey Agar, Blood Agar and Chocolate Agar media⁴ with 7% sheep blood. Isolates obtained in significant numbers i.e., 10^5 cfu/ml of specimen were then identified following Bergey's Manual of Systematic Bacteriology. β -haemolytic Streptococci were

grouped using commercially available Oxoid Streptococcal Grouping kit. *Invitro* antimicrobial susceptibility testing of the isolates were done using Kirby-Bauer⁵ Disc Diffusion Technique and zone size was interpreted following NCCLS⁶ standard. All the plates were incubated at 37°C and read after 24 hours incubation. Also, Giemsa staining was done for *Pneumocystis carinii* and for fungi KOH preparation was performed. Likewise, saline preparation was performed for *Paragonimus westermanii*.⁷

Results

Out of total 181 samples processed only 75 (41.4%) had microbiological diagnosis, 102 (56.4%) gave non-significant growth and remaining 4 (2.2%) gave no growth (Fig. 1).




Fig. 1: Pattern of growth

Out of total 87 bacterial isolates, 63 (72.4%) were Gram-negative and 24 (27.6%) Gram-positive as depicted in (Fig. 2).





Fig. 2: Distribution of Bacterial Isolates

Among the isolates *H. influenzae* was most frequently encountered at 20 (23.0%). *S pneumoniae* was most common Gram-positive isolates accounting for 11 (12.7%) (Fig. 3).


Table 1: Prevalence of LRTI by age and gender

| Age groups (Years) | Male | | Female | | Total positive cases No. % |
|--------------------|----------------------|----------------|----------------------|----------------|----------------------------|
| | No. of cases studied | Positive cases | No. of cases studied | Positive cases | |
| >14-19 | 7 | 5 55.5% | 9 | 4 44.5% | 9 12.0% |
| 20-29 | 22 | 9 75% | 16 | 3 25% | 12 16% |
| 30-39 | 13 | 8 57.1% | 10 | 6 42.9% | 14 18.7% |
| 40-49 | 10 | 5 55.5% | 8 | 4 44.5% | 9 12.0% |
| 50-59 | 24 | 10 62.5% | 18 | 6 37.5% | 16 21.3% |
| 60-69 | 19 | 8 80% | 12 | 2 20% | 10 13.3% |
| 70-79 | 5 | 1 33.3% | 4 | 2 66.7% | 3 4.0% |
| 80-89 | 2 | 0 0% | 2 | 2 100% | 2 2.7% |
| Total | 102 | 46 61.3% | 79 | 29 38.7% | 75 41.4% |


**Fig. 3:** Bacterial involvement in LRTI (In Number)

LRTI was found to be more prevalent in males than females. Of the total 102 males and 79 females enrolled for the study, LRTI was established in 61.3% and 38.7% respectively. Among the LRTI infected cases, highest prevalence was detected in age group 50-59 yrs followed by 30-39 yrs as depicted in (Table 1).


Among the different ethnic groups, LRTI was found highly prevalent in Nepali/Pariyar/Kumal (62.5%) followed by Gurung/Magar/Rai/Tamang (46.5%) and least prevalent in Kshetrys (23.5%) as shown in (Fig. 4).

**Fig. 4:** Distribution of LRTI among various ethnic groups

In vitro antibiotic susceptibility test showed that for Gram-positive bacteria Ciprofloxacin at 79.2% was the most effective antibiotic and Cotrimoxazole least effective (Fig. 5).

**Fig. 5:** Antibiotics susceptibility Pattern of Gram Positive bacteria

Similarly, for Gram-negative isolates Chloramphenicol (100%) was the most effective antibiotic followed by Amikacin (79.1%) and Ciprofloxacin (66.7%). Cotrimoxazole was the least effective of all (Fig. 6).

**Fig. 6:** Antibiotics susceptibility Pattern of Gram Negative Bacteria

Discussion

Based on the laboratory findings (i.e., significant growth obtained in culture) LRTI was found prevalent in only 75 (41.1%) of the total 181 cases investigated. In a similar study carried out by Hosker⁸ (1994) significant growth was obtained in about 60% of the cases. The greater negative results in the current study may be attributed to viral and

Lower respiratory tract infection in adults

aetiological agents, apart from bacteria. Though the study also attempted to look for *Pneumocystis carinii* and *Paragonimus westermanii*; no such findings were obtained.

Altogether 15 different types of bacteria were isolated and identified in sputum culture. Greater number of the isolates was GNB. The ratio of Gram-negative bacterial infection to Gram-positive bacterial infection was 2.6:1. These findings correlated well with that of similar study conducted in 1995.⁹

The study showed males (61.3%) at more risk to LRTI than females (38.7%). Reason as such could be because females enrolled in the study comprised largely of housewives, who, being less mobile experienced less exposure to respiratory risk factors. It could also be attributed to predisposing factors in males like smoking and alcoholism.¹⁰ Furthermore, in this study higher prevalence of LRTI was observed in elderly population (40 years and older) accounting for 53.3% of the total infected cases than younger lot (30 years and younger at 46.7%). In particular, patients in age groups 50-59 years were found more prone to LRTI (21.3%) and the prevalence decreased drastically in population 80 years and older (2.7%). Similar study showed that pneumococcal infection is common in patients over the age of 55, who also have an increasing incidence of LRTI as they get older¹¹, since immune system in older patients become less effective owing to either malnutrition or underlying degenerative diseases such as diabetes mellitus, emphysema, uraemia etc.¹²

The current study also indicated plausible link between LRTI and ethnic group enrolled. Accordingly, Kumal/Nepali/Pariyar (62.5%) was found more susceptible to LRTI and Kshetrys were least susceptible of all at 23.5%. This discrepancy in distribution of LRTI among various ethnic groups could be attributed to their social, cultural and religious background not mentioning the inequality in access to services and in the range of opportunities available; however, the study did not divulge the precise link between the prevalence of LRTI and ethnic groups.

Of the total bacterial isolates *Haemophilus influenzae* was the most common isolates (23%) followed by *klebsiella* (18.3%) and *pseudomonas* (14%). *Streptococcus pneumoniae* (12.3%) was the predominant pathogen among the Gram-positive isolates. Though earlier study had shown *S. pneumoniae* as a predominant isolate followed by *H. influenzae*¹¹, our findings correlates well with that of Sayami⁹, 1994. It is also noteworthy that in the current study *H. influenzae* (and other GNB) were diagnosed predominantly in population with pre-existing chronic diseases.¹¹

In this study isolates resistant to one or more antibiotics commonly in use was observed to be higher than similar

study conducted in 1999.¹³ All Gram-positive isolates showed resistance to Cotrimoxazole whereas 79.4% of Gram-negative isolates were resistant to Cotrimoxazole. Similarly, higher percentage of both Gram-positive and Gram-negative isolates showed resistant to Ciprofloxacin, Cephalexin and Ampicillin compared to earlier study.

Pseudomonas and *Klebsiella* are among the bacteria manifesting high levels of resistance-most notably in developing nations. The increase in the antibiotic non-susceptible strains of pathogens in recent years could be attributed to their indiscriminate and promiscuous use. This alarming situation is due, in part, to widespread confusion over the difference between viral and bacterial respiratory infections which has also led to the emergence of resistant pneumococci³ and staphylococci. Similarly, in case of GPB like *S. pneumoniae* the use of suboptimal and long duration regimens increase the opportunity for acquisition and/or amplification of resistant strains.¹⁵ Formerly, first-line medications were both effective and affordable. With the onset of resistance however, newer treatments are proving too costly to the vast majority of those living in poor developing nations.³

Conclusion

The study indicates that there has not been any significant changes in the pattern of bacterial involvement in causing LRTI. Also, the population and the age groups at risk have remained more or less the same over the period of time. However, increasing number of drug resistant pathogens emphasizes the importance of judicious antibiotic use. Educational program in this respect for general public and clinician as well could be useful in checking the emergence of drug resistant pathogens. Furthermore, including atypical pathogens like *Mycoplasma pneumonia* and other respiratory viruses in microbial investigation of LRTI could reduce the unnecessary antibiotic use in humans.

Acknowledgements

We would like to extend out heartfelt gratitude to Prof. Nhuchhe Ratna Tuladhar, Head, Department of Clinical Microbiology, Institute of Medicine, Tribhuvan University Teaching Hospital, Maharajgunj, Kathmandu Nepal and Mr. Govinda Gurung, Health Research Laboratory, Institute of Medicine for their kind support and co-operation.

References

1. World Health Organization. Disease Burden and economics. *Practical Approach to Lung Health (PAL)*. WHO 2003.

2. Collee JG and Watt B. "Bacterial infection of respiratory tract". In Topley and Wilson's Principles of Bacteriology, Virology, and Immunity Vol II, 8th edition. BC Decker, Inc. UK 1990.
3. World Health Organization. The Big Guns of Resistance. Infectious disease control. WHO 2000.
4. Collee JG, Fraser AG, Dugid JG and Marmion BP. Mackie & Mc Cartney Practical Medical Microbiology, 14th edition. Churchill Livingstone, UK 1996.
5. Bauer AW, Kirby W MM, Sherris JC. Antibiotic susceptibility testing by a standardized single disc method. *Am J Clin Path* 1966; **45**: 493-496.
6. National Committee for Clinical Laboratory Standards (NCCLS). Performance standards for antimicrobial disc susceptibility tests; Approved Standards, 7th edition. 2000.
7. Cheesbrough M. *District Laboratory Practice in Tropical Countries Part II*. Low price edition. Cambridge University Press, UK 2002.
8. Hosker H. Management of Community Acquired LRTI. *BMJ* 1994; **308**: 871-872.
9. Sayami A, Shrestha B. Critical Care: *Manual of ICU and CCU*. Tribhuvan University Teaching Hospital 1995.
10. Doddannavar RP. Changing profile of Bacterial Pneumonia. *Journal of Indian Medical Association* 1985; **83**: 149-153.
11. MacFarlane JT, Colville A, Guion A, Macfarlane RM, Rose DH. Prospective study of aetiology and outcome of adult lower-respiratory-tract infections in the community. *Lancet* 1993; **341**: 511-514.
12. Berk LS. Bacterial infections in the elderly. *Post Graduate Medicine* 1985; **77**: 168-173.
13. Joshi K, Basnet SR, Pokhrel BM and Tuladhar NR. A prospective study on bacteriology of Lower Respiratory Tract Infection among the patients visiting TUTH Kathmandu (1999).
14. Schrag JS., Beall B, Dowell S. Resistant pneumococcal infections: the burden of disease and challenges in monitoring and controlling antimicrobial resistance. WHO 2001.